

LINEAR AND QUADRATIC CHABAUTY

FOR AFFINE HYPERBOLIC CURVES

by Marius Leonhardt, Steffen Müller

 X/\mathbb{Q} smooth proj. curve of genus $g \geq 2$ p prime of good red., $\mathcal{U}^{\text{ét}} \rightarrowtail \mathcal{U}$ $G_{\mathbb{Q}}$ -equiv. quotient

Chabauty - Kim diagram

$$\begin{array}{ccc} X(\mathbb{Q}) & \longrightarrow & X(\mathbb{Q}_p) \\ \downarrow j & & \downarrow j_p \\ \text{Sel}_{\mathcal{U}}(X) & \xrightarrow{\text{loc}_p} & H^1_f(G_p, \mathcal{U}) \end{array}$$

Chabauty - Kim locus

$$X(\mathbb{Q}_p)_u := \{ x \in X(\mathbb{Q}_p) : j_p(x) \in \text{loc}_p(\text{Sel}_u(X)) \}$$

$$X(\mathbb{Q}) \subseteq X(\mathbb{Q}_p)_u \subseteq X(\mathbb{Q}_p)$$

Conj: $\# X(\mathbb{Q}_p)_u < \infty$ for u sufficiently large

- Linear Chabauty: $\mathcal{U} = \mathcal{U}_1^{\text{ét}} = (\mathcal{U}^{\text{ét}})^{\text{ab}} \rightarrow X(\mathbb{Q}_p)_1$

Thm: Let $r_p := \text{rk}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(\text{Jac}_X)$ ($= r$ under Tate-Shafarevich conj.).If $g - r_p > 0$, then $\# X(\mathbb{Q}_p)_1 < \infty$.

- Quadratic Chabauty: $\mathcal{U} = \mathcal{U}_{\text{QC}} = \text{certain intermediate quotient } \mathcal{U}_2^{\text{ét}} \rightarrowtail \mathcal{U}_{\text{QC}} \rightarrow \mathcal{U}_1^{\text{ét}}$
 $\rightarrow X(\mathbb{Q}_p)_{\text{QC}}$

Thm: (Balakrishnan - Dogra)Let $g_f := \text{rk NS}(\text{Jac}_X) + \text{rk NS}(\text{Jac}_{X_{\bar{\mathbb{Q}}}})^{\sigma = -1}$, $\sigma = \text{complex conj.}$ If $g + g_f - 1 - r_p > 0$, then $\# X(\mathbb{Q}_p)_{\text{QC}} < \infty$.

Aim: generalise this to affine hyperbolic curves

Setup: Y/\mathbb{Q} affine hyperbolic curve, $Y = X \setminus D$ with X projective, $n := \#D(\bar{\mathbb{Q}})$

- Ex:
- $Y = \mathbb{P}^1 \setminus \{0, 1, \infty\}$,
 - $Y = E \setminus \{O\}$ with E elliptic curve,
 - $Y: y^2 = f(x)$ affine hyperell. curve

S finite set of primes

$\mathbb{Z}_S = \{x \in \mathbb{Q} : v_l(x) \geq 0 \quad \forall l \notin S\}$ ring of S -integers

$Y = X \setminus D$ regular model of $Y = X \setminus D$ over \mathbb{Z}_S

Thm: (Siegel, Faltings)

$$\#Y(\mathbb{Z}_S) < \infty.$$

Variant of Chabauty-Kim for S -integral points:

$p \notin S$ s.t. X_{F_p} and D_{F_p} smooth, choose base point in $Y(\mathbb{Z}_S)$ for $U^{\text{ét}}$,

$U^{\text{ét}} \rightarrow U$ $G_{\mathbb{Q}}$ -equiv. quotient.

Chabauty-Kim diagram:

$$\begin{array}{ccc} Y(\mathbb{Z}_S) & \longrightarrow & Y(\mathbb{Z}_p) \\ \downarrow j & & \downarrow j_p \\ \text{Sel}_{S,u}(Y) & \xrightarrow{\text{loc}_p} & H_f^1(G_p, U) \end{array}$$

Chabauty-Kim locus:

$$Y(\mathbb{Z}_p)_{S,u} := \{y \in Y(\mathbb{Z}_p) : j_p(y) \in \text{loc}_p(\text{Sel}_{S,u}(Y))\}$$

- Linear Chabauty: $U = U_i^{\text{ét}} \rightarrow Y(\mathbb{Z}_p)_{S,1}$
- Quadratic Chabauty: construct suitable intermediate quotient
 $U_i^{\text{ét}} \rightarrow U_{QC} \rightarrow U_i^{\text{ét}} \rightarrow Y(\mathbb{Z}_p)_{S,QC}$

Invariants attached to (Y, S, ρ) :

- $r_p = \text{rk}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(\text{Jac } X)$ p^∞ -Selmer rank
- $g = \text{genus of } X$
- $\beta_f = \text{rk } NS(\text{Jac } X) + \text{rk } NS(\text{Jac } X_{\bar{\mathbb{Q}}})^{\sigma = -1}$
- $n = \# D(\bar{\mathbb{Q}})$ number of geom. pts at infinity
 $= n_1 + 2n_2$ with $n_1 = \# D(R)$, $2n_2 = \#(D(C) \setminus D(R))$
- $b := \#|D| + n_2 - 1 \quad (\geq 0 \text{ since } Y \text{ affine})$
- $s := \#S$

Theorem A: (Leonhardt - L. - Müller)

- (1) If $\alpha_1 := g - r_p + b - s > 0$, then $\# Y(\mathbb{Z}_p)_{S,1} < \infty$.
- (2) If $\alpha_2 := \alpha_1 + \beta_f > 0$, then $\# Y(\mathbb{Z}_p)_{S, \text{QC}} < \infty$.

Theorem B: (LLM)

- (1) If $\beta_1 := \frac{1}{2}g(g+3) - \frac{1}{2}\gamma(r+3) + b - s > 0$, then
 $\# Y(\mathbb{Z}_p)_{S,1} \leq K_p \cdot \prod_{l \in S} (n_l + n) \cdot \prod_{l \notin S} n_l \cdot \# Y(\mathbb{F}_p) \cdot (4g + 2n - 2)^2 (g+1)$
- (2) If $\beta_2 := \beta_1 + \beta_f > 0$, same bound for $Y(\mathbb{Z}_p)_{S, \text{QC}}$.

Here: $K_p := 1 + \frac{p-1}{(p-2)\log(p)}$ (p odd), $K_2 := 2 + \frac{2}{\log(2)}$.

$n_l := \#\text{irred. cpts of the mod-}l \text{ special fibre of } X_{\mathbb{F}_p}$ (if $l \notin S$)

resp. of the minimal regular normal crossings model of (X, D)
over \mathbb{Z}_l (if $l \in S$)

Ranks: • Have similar results for $Y(\mathbb{Z}_p)_{S, \text{wt} \geq 2}$, $U_2^{\text{ét}} \rightarrow U_{\text{wt} \geq 2} \rightarrow U_{\text{QC}} \rightarrow U_1^{\text{ét}}$.

$$h_{BK} := \dim_{\mathbb{Q}_p} H^1_f(G_{\mathbb{Q}}, \text{Hom}(A^2 V_p \text{Jac } X, \mathbb{Q}_p(1)))$$

(= 0 conjecturally by Bloch - Kato)

C. If $g^2 - r_p + g + b - s > 0$, then $\#Y(\mathbb{Z}_p)_{S, \text{wt} \geq -2} < \infty$

D. If $\frac{1}{2}g(3g+1) - \frac{1}{2}r_p(r_p+3) + g + b - s - h_{BK} > 0$,

then $Y(\mathbb{Z}_p)_{S, \text{wt} \geq -2}$ satisfies bound from Thm B.

- "Balakrishnan-Dogra trick" \rightarrow CK loci $Y(\mathbb{Z}_S)_{S, u}^{\text{BD}}$ where Theorems hold with r instead of r_p .
- Bettis-Corwin-Leonhardt: Bound on $Y(\mathbb{Z}_p)_{S, \infty}$ assuming Tate-Shafarevich + Bloch-Kato.

1. Refined Selmer schemes

For all ℓ have diagram

$$\begin{array}{ccc} Y(\mathbb{Z}_S) & \longrightarrow & Y(Q_\ell) \\ \downarrow j & & \downarrow j_\ell \\ H^1(G_Q, U) & \xrightarrow{\text{loc}_\ell} & H^1(G_\ell, U) \end{array} \quad \begin{array}{l} \text{can replace } Y(Q_\ell) \text{ with} \\ Y(\mathbb{Z}_\ell) \text{ for } \ell \notin S \end{array}$$

Selmer functor: $R \mapsto \{\alpha \in H^1(G_\ell, U) : \text{loc}_\ell(\alpha) \in \begin{cases} j_\ell(Y(\mathbb{Z}_\ell))^{\text{zar}}, & \ell \notin S, \\ j_\ell(Y(Q_\ell))^{\text{zar}}, & \ell \in S. \end{cases}\}$

This is representable by the (refined) Selmer scheme $\text{Sel}_{S, u}(y)$.
 \rightarrow CK diagram

Say $x, y \in Y(\mathbb{Z}_S)$ have same reduction type if

$\forall \ell$: mod- ℓ reductions on the same irr. cpt. or (if $\ell \in S$)
 the same cuspidal point.

reduction type: $\Sigma = (\Sigma_\ell)_{\ell \text{ prime}}$

Selmer scheme is union $\text{Sel}_{S, u}(y) = \bigcup_{\Sigma} \text{Sel}_{\Sigma, u}(y)$,

corresponding to $Y(\mathbb{Z}_S) = \coprod_{\Sigma} Y(\mathbb{Z}_S)_{\Sigma} \leftarrow \text{points of red. type } \Sigma$

$$\rightarrow \mathcal{Y}(\mathbb{Z}_p)_{S, u} = \bigcup_{\Sigma} \mathcal{Y}(\mathbb{Z}_p)_{S, u, \Sigma}$$

Σ -refined CK diagram

$$\begin{array}{ccc} \mathcal{Y}(\mathbb{Z}_S)_\Sigma & \hookrightarrow & \mathcal{Y}(\mathbb{Z}_p) \\ \downarrow j & & \downarrow j_p \\ \text{Sel}_{\Sigma, u}(\mathcal{Y}) & \xrightarrow{\text{loc}_p} & H_f^1(G_p, U) \end{array}$$

Strategy for finiteness of CK loci:

$$\underline{\text{If}} \dim \text{Sel}_{\Sigma, u}(\mathcal{Y}) < \dim H_f^1(G_p, U)$$

$\Rightarrow \text{loc}_p$ not dominant

$$\Rightarrow \exists 0 \neq f \text{ s.t. } f \circ \text{loc}_p = 0$$

$$\Rightarrow \mathcal{Y}(\mathbb{Z}_p)_{S, u, \Sigma} \subseteq V(f \circ j_p) \text{ finite}$$

Can compute dimensions using (abelian) Galois cohomology.

2. Weight filtrations on Selmer schemes

Betti's filtration $\dots \subseteq W_{-2}U \subseteq W_{-1}U = U$ by subgroup schemes

$$\text{s.t. } [W_{-i}U, W_{-j}U] \subseteq W_{-(i+j)}U \quad \forall i, j \geq 1.$$

$$\Rightarrow \text{gr}_{-k}^W U := W_{-k}U/W_{-k-1}U \text{ rep'n of } G_{\mathbb{Q}} \text{ on fin. dim. } \mathbb{Q}_p\text{-v.s.}$$

$$\text{Sel}_{\Sigma, u} \hookrightarrow \prod_{k=1}^{\infty} H_f^1(G_{\mathbb{Q}}, \text{gr}_{-k}^W U) \times \prod_{\ell \in S} \bigoplus_{\substack{\text{crystalline at } p, \\ \text{unram. away from } p}} \mathcal{G}_{\Sigma_{\ell}} \quad (\text{non-canonically})$$

$$\Rightarrow \dim \text{Sel}_{\Sigma, u} \leq s + \sum_{k=1}^{\infty} \dim_{\mathbb{Q}_p} H_f^1(G_{\mathbb{Q}}, \text{gr}_{-k}^W U),$$

$$H_f^1(G_p, U) \cong \prod_{k=1}^{\infty} H_f^1(G_p, \text{gr}_{-k}^W) \quad (\text{non-canonically})$$

$$\Rightarrow \dim H_f^1(G_p, U) = \sum_{k=1}^{\infty} \dim_{\mathbb{Q}_p} H_f^1(G_p, \text{gr}_{-k}^W)$$

- Linear Chabauty: $\mathcal{U} = \mathcal{U}_1^{\text{ét}} = \mathcal{U}_Y^{\text{ab}}$

$Y \hookrightarrow X$ induces $\mathcal{U}_Y^{\text{ab}} \rightarrow \mathcal{U}_X^{\text{ab}} = V_p \mathbb{Z}_p \text{Jac}_X := \left(\varprojlim_n \mathbb{Z}_p \text{Jac}_X(\bar{\mathbb{Q}})[p^n] \right) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$

$$1 \rightarrow \underbrace{\mathbb{Q}_p(1)^{\mathcal{D}(\bar{\mathbb{Q}})}}_{gr_{-2}^w =: I} / \mathbb{Q}_p(1) \rightarrow \mathcal{U}_Y^{\text{ab}} \rightarrow \underbrace{V_p \mathbb{Z}_p \text{Jac}_X}_{gr_{-1}^w} \rightarrow 1$$

Can compute local and global Galois coh:

$$\text{wt -1: } \dim H_f^1(G_{\mathbb{Q}}, V_p \mathbb{Z}_p \text{Jac}_X) = r_p$$

$$\dim H_f^1(G_p, V_p \mathbb{Z}_p \text{Jac}_X) = g$$

$$\text{wt -2: } \dim H_f^1(G_{\mathbb{Q}}, I) = n_1 + n_2 - |\mathcal{D}|$$

$$\dim H_f^1(G_p, I) = n - 1$$

If $\alpha_1 > 0$ in Thm A

$$\Rightarrow \dim \text{Sel}_{\Sigma, \mathcal{U}} < \dim H_f^1(G_p, \mathcal{U})$$

$$\Rightarrow \# \mathcal{U}(\mathbb{Z}_p)_{S, 1} < \infty$$

- Quadratic Chabauty: Construction of \mathcal{U}_{QC}

$$1 \rightarrow \Lambda^2 V_p \mathbb{Z}_p \text{Jac}_X \oplus I \rightarrow \mathcal{U}_Y / W_{-3} \mathcal{U}_Y \rightarrow V_p \mathbb{Z}_p \text{Jac}_X \rightarrow 1$$

\downarrow max. Artin-Tate quotient \downarrow ||

$$1 \rightarrow \left(\mathbb{Q}_p \otimes_{NS(\text{Jac}_X(\bar{\mathbb{Q}}))} \mathcal{D}(1) \right) \oplus I \rightarrow \mathcal{U}_{QC} \rightarrow V_p \mathbb{Z}_p \text{Jac}_X \rightarrow 1$$

η

can compute its Galois coh.

3. Bounding CK loci

Weight filtration on $\mathcal{O}(\text{Sel}_{\Sigma, \mathcal{U}})$ and $\mathcal{O}(H_f^1(G_p, \mathcal{U}))$ by fin. dim. subspaces, preserved by $\text{loc}_p^{\#}: \mathcal{O}(H_f^1) \rightarrow \mathcal{O}(\text{Sel}_{\Sigma, \mathcal{U}})$

Assume $\dim W_m \mathcal{O}(\text{Sel}_{\Sigma, \mathcal{U}}) < \dim W_m \mathcal{O}(H_f^1)$

$$\Rightarrow \exists 0 \neq f \in W_m \mathcal{O}(H_f^1) \text{ s.t. } \text{loc}_p^{\#} f = 0$$

$\Rightarrow f \circ j_p : Y(\mathbb{Z}_p) \rightarrow \mathbb{Q}_p$ is "Coleman algebraic function of weight $\leq m$ "

Betts: bound on number of zeroes in each residue disc

\Rightarrow get bound on $Y(\mathbb{Z}_p)_{S, U, \Sigma}$ and $Y(\mathbb{Z}_p)_{S, U}$

In our theorems we get weight 2 functions

\rightarrow sum of double and single Coleman integrals and rat. functions

Compute $\dim W_m \mathcal{O}(Z)$ via Hilbert series.

$$HS_Z(t) := \sum_{i=0}^{\infty} \dim \text{gr}_i^W \mathcal{O}(Z) t^i \in \mathbb{N}_0[[t]]$$

$$\text{Betts: } HS_{\text{Sel}_{\Sigma, U}}(t) \lesssim (1-t^2)^{-s} \prod_{k=1}^{\infty} (1-t^k)^{-\dim H_f^1(G_Q, \text{gr}_{-k}^W U)} =: HS_{\text{glob}}(t)$$

$$HS_{H_f^1}(t) = \prod_{k=1}^{\infty} (1-t^k)^{-\dim H_f^1(G_p, \text{gr}_{-k}^W U)} =: HS_{\text{loc}}(t)$$

For $U = U_1 e^{st}$:

$$HS_{\text{glob}}(t) = 1 + r_p t + (s + n_1 + n_2 - \#|D| + \frac{1}{2} r_p(r_p+1)) t^2 + \dots$$

$$HS_{\text{loc}}(t) = 1 + g t + (n-1 + \frac{1}{2} g(g+1)) t^2 + \dots$$

If $n_1 > 0$ in Thm B

$\Rightarrow t^2\text{-coeff of } HS_{\text{glob}}(t) < t^2\text{-coeff of } HS_{\text{loc}}(t)$

$\Rightarrow \exists$ nonzero Coleman alg. function $Y(\mathbb{Z}_p) \rightarrow \mathbb{Q}_p$ of weight ≤ 2

vanishing on $Y(\mathbb{Z}_p)_{S, 1, \Sigma}$

\Rightarrow bound on $\# Y(\mathbb{Z}_p)_{S, 1, \Sigma}$ and $\# Y(\mathbb{Z}_p)_{S, 1}$